

PLEUROTUS OSTREATUS (JACQ.) P. KUMM. E PLEUROTUS ERYNGII (F.) QUÉL. COMO POTENCIAIS AGENTES MICORREMEDIADORES DE EFLUENTES CONTAMINADOS COM BENTAZONA E 2,4-D

Pleurotus ostreatus (Jacq.) P. Kumm. and *Pleurotus eryngii* (F.) Qué. as potential mycoremediator agents of wastewater contaminated with bentazon and 2,4-D

Graciela Rozza^{1,2}; Kézia Melo²; Thaynã Gonçalves Timm¹; Judith Paola Urón Santiago³; Admir José Giachini³; Jaqueline da Silva Coelho-Moreira⁴; Maria Pilar Serbent^{2*}

Resumo: Micorremediação tem sido estudada como alternativa para remoção de pesticidas de efluentes. Foi avaliada a capacidade *in vitro* dos fungos basidiomicetos *Pleurotus ostreatus* e *Pleurotus eryngii* como potenciais agentes de remediação de efluentes que contêm herbicidas. *P. ostreatus* e *P. eryngii* foram cultivados em meios líquidos contendo, respectivamente, bentazona (4,5 g L⁻¹) e 2,4-D (5,4 g L⁻¹), analisando a capacidade de crescimento dos fungos (biomassa seca), remoção dos compostos (CG/MS) e toxicidade do meio após tratamento com os fungos (testes *Allium cepa*). *P. ostreatus* e *P. eryngii* se mostraram tolerantes à bentazona e 2,4-D, respectivamente, uma vez que cresceram nos meios líquidos. A análise CG/MS mostrou que os fungos não foram eficientes na remoção da bentazona e do 2,4-D após 7 e 21 dias de cultivo, respectivamente. Em ambos os casos os efluentes finais resultaram tóxicos para *A. cepa*. Apesar de não ter sido comprovada a remoção de pesticidas, os resultados do trabalho mostram que *P. ostreatus* e *P. eryngii* têm o potencial de tolerar a bentazona e o 2,4-D em meios líquidos. Portanto, algumas condições de cultivo diferentes devem ser testadas para explorar o potencial destes fungos para a remoção de pesticidas.

Abstract: Mycoremediation has been studied as an alternative to removing pesticides from wastewater. Here, we evaluated the *in vitro* capability of the basidiomycetes *Pleurotus ostreatus* and *Pleurotus eryngii* as potential remediation agents of effluents containing the pesticides bentazon and 2,4-D, respectively. *P. ostreatus* and *P. eryngii* were cultivated in liquid media containing bentazon (4.5 g L⁻¹) and 2,4-D (5.4 g L⁻¹), respectively. Biomass dry weight and CG/MS analysis were performed to evaluate the fungi' capability to grow in the contaminated media as well as to degrade the studied pesticides. To evaluate the toxicity of the wastewater after the cultivation of the fungi, tests with *Allium cepa* were conducted. *P. ostreatus* and *P. eryngii* presented tolerance to the bentazon and 2,4-D, respectively. CG/MS analysis showed that the fungi were not efficient in the removal of bentazon and 2,4-D after seven and twenty-one days of cultivation, respectively. Finally, in both cases, the final effluents were toxic to *A. cepa*. Despite the non-removal of pesticides, *P. ostreatus* and *P. eryngii* have the potential to tolerate bentazon and 2,4-D in liquid media. Therefore, some different cultivation conditions must be tested to explore the potential of these fungi for pesticide removal.

Palavras-chave:

Saneamento; Águas residuais; Pesticidas; Basidiomicetos; Micorremediação.

Keywords:

Sanitation; Wastewater; Pesticides; Basidiomycetes; Mycoremediation.

¹Regional University of Blumenau, Environmental Engineering Graduate Program, Blumenau, Santa Catarina, Brazil. rozzagraciela@gmail.com; ttimm@furb.br

²Santa Catarina State University, Department of Sanitary Engineering, Ibirama, Santa Catarina, Brazil. keziameo95@gmail.com; mariapilar.serbent@udesc.br

³Federal University of Santa Catarina, Department of Microbiology, Immunology, and Parasitology, Florianópolis, Santa Catarina, Brazil. paolauron@gmail.com; admir.giachini@gmail.com

⁴Federal Institute of Paraná, Ivaiporã, Paraná, Brazil. jaquecoelho@hotmail.com

1 INTRODUCTION

Pesticides are chemical compounds widely used to protect crops from undesirable organisms, such as weeds, fungi, insects, and pathogens (SJERPS et al., 2019). Nevertheless, several of these pesticides remain in the effluents of conventional crops, which are normally discharged untreated into waterways (YADAV et al., 2015). Consequently, the use of these substances worldwide has led to the contamination of environmental matrices as ground and surface water, including drinking water (SJERPS et al., 2019; QUINTANA et al., 2019; CHEN et al., 2018; XIE et al., 2019; XU et al., 2019; CALDAS et al., 2013; POSTIGO et al., 2010; KÖCK-SCHULMEYER et al., 2014; VIEIRA et al., 2016). It is a major environmental and public health problem since most of these compounds offer a high-level risk to aquatic organisms. Therefore, the occurrence of pesticides in the environment has been extensively studied in the last decades (CHEN et al., 2018; XIE et al., 2019).

Among the pesticides that may remain in wastewater, there are the 3-isopropyl-1H-2,1,3-benzothiadiazin-4(3H)-one-2,2-dioxide (Bentazon) and the 2,4-dichloro phenoxy acetic acid (2,4-D). Bentazon is a post-emerging herbicide largely applied in cereal crops and it is one of the most widely used herbicides in conventional agriculture and gardening (SALMAN; HAMEED, 2010). Besides, the systemic herbicide 2,4-D is used globally in the selective control of nest plants in agriculture and forestry (ZUANAZZI et al., 2020). Several studies have related the occurrence of bentazon in the environment (CALDAS et al., 2013; POSTIGO et al., 2010; VIEIRA et al., 2016) and recent studies have shown the contamination of surface and underground water bodies by 2,4-D (ECHEVERRI GONZÁLEZ et al., 2019; CASTRO LIMA et al., 2020).

The findings of the occurrence and risks of pesticides have heightened the need for advanced treatments to remove the contaminants from the environment since conventional treatment is not capable to remove such compounds. In this context, mycoremediation has been studied as an economical, eco-friendly, and effective biological tool that recruits fungi to degrade, transform or immobilize contaminants from the environment (PURNOMO et al., 2010; COELHO-MOREIRA et al., 2018; PEREIRA et al., 2013; CUPUL et al., 2014; FREITAS et al., 2017; GANASH et al., 2016; SERBENT et al., 2019; 2020; KAPAH; SACHDEVA, 2017; BOSCO; MOLLEA, 2019; AKHTAR; MANNAN, 2020).

Fungi tend to degrade a vast number of contaminants (HARMS et al., 2011; SINGH, 2006; VALDEZ-VAZQUEZ et al., 2020; ZAPANA-HUARACHE et al., 2020) and the white-rot fungi (WRF) are one of the major ecological categories of wood-decaying fungi (RILEY et al., 2014) responsible for complex ecosystem processes going on during wood decay and nutrient cycling (LEONHARDT et al., 2019). WRF can be promising candidates for the treatment of contaminated wastes (AKHTAR; MANNAN, 2020), but their potential in bioremediation processes regarding chlorinated compounds is still far from being fully explored (SERBENT et al., 2020). Furthermore, basidiomycetes, such as the *Pleurotus* genre, are known for producing enzymes including lignin peroxidase, manganese peroxidase, and laccase, which promote the degradation of substances (PURNOMO et al., 2010; SINGH, 2006; ZAPANA-HUARACHE et al., 2020).

In this context, this study aims to evaluate the *in vitro* tolerance and growth of *Pleurotus ostreatus* (Jacq.) P. Kumm. and *Pleurotus eryngii* (Fr.) Quél. in liquid medium contaminated with bentazon and 2,4-D, respectively, as well as the capability of these fungi to remove the corresponding herbicide from spiked water and evaluation of its toxicity by *Allium cepa* test after treatments.

2 MATERIAL AND METHODS

2.1 Source of pesticides and fungi


Bentazon was obtained from its commercial form, Basagram®, containing 600 g L⁻¹ of the active ingredient, and 2,4-D was obtained from its commercial form, Aminol®, containing 670 g L⁻¹ of the active ingredient. In both cases, the spiked water was prepared with sterilized and deionized water. *P. ostreatus* and *P. eryngii* were provided by the culture collection of the laboratory of microorganisms and biotechnological processes at the Federal University of Santa Catarina.

2.2 In vitro evaluation of pesticides tolerance and removal by strains of *Pleurotus*

P. ostreatus and *P. eryngii* were cultivated in Petri dishes containing potato dextrose agar (PDA) at 27 ± 2 °C before inoculation in the liquid medium and its subsequent analyses.

Figure 1 summarizes the studied treatments and each analysis for the *in vitro* evaluation of pesticide tolerance and removal by *Pleurotus* fungi.

Figure 1. Studied treatments and each analysis for the *in vitro* evaluation of pesticide tolerance and removal by *Pleurotus* fungi.



2.3 Fungal growth in liquid culture media

To establish the tolerance and removal of bentazon by *P. ostreatus* and 2,4-D by *P. eryngii*, experiments of liquid cultures were conducted in a stationary state.

To represent real concentrations of the pesticides in wastewater (CUPUL et al., 2014; FREITAS et al., 2017; GANASH et al., 2016; SERBENT et al., 2019; 2020), the bentazon and 2,4-D were diluted in deionized water separately, resulting in 4.5 g L⁻¹ (T0_{BZ}) and 5.4 g L⁻¹ (T0_{2,4D}) initial solutions, respectively. Deionized water was used as a control treatment for both pesticide experiments. Each sample was prepared in a glass flask previously sterilized.

For bentazon experiments, five mycelial disks (diameter: 8.0 mm) of *P. ostreatus* were inoculated in T0_{BZ} (T1_{BZ}) and control treatment (CT_{BZ}). The samples (T0_{BZ}, T1_{BZ}, and CT_{BZ}) were incubated at 27 ± 1 °C for 7 days (SANTOS et al., 2019; KIST et al., 2020).

For 2,4-D experiments, five mycelial disks (diameter: 8.0 mm) of *P. eryngii* were inoculated in T0_{2,4D} (T1_{2,4D}) and control treatment (CT_{2,4D}). The samples (T0_{2,4D}, T1_{2,4D}, and CT_{2,4D}) were incubated at 27 ± 1 °C for 21 days.

2.4 Biomass dry weight of fungi and pesticides concentration

After the incubation period, the biomass dry weight of the *P. ostreatus* and *P. eryngii* in liquid cultures were performed according to the procedure proposed by Souza et al. (2006).

Bentazon and 2,4-D concentration in the samples was analyzed by gas chromatography/mass spectrometry (GC/MS) as determined by the EPA 8270D method (US EPA, 2014), in which the quantification limit is 0.1 µg L⁻¹.

2.5 Toxicity evaluation of treated water by *Allium* test

The toxicity test was conducted with *Allium cepa* L. (onion) samples as the adapted method used by Marinho et al. (2017). First, 20 onions with similar shapes and sizes were selected, peeled, and their roots were carefully removed. Then, the onions were placed in plastic cups with tap water in the dark for 72 hours. After this period, the roots were observed and the ones with similar sizes were selected for the toxicity test. These roots were removed, and the bulbs were placed in plastic cups with the samples T0_{BZ}, T1_{BZ}, T0_{2,4D}, and T1_{2,4D} in the dark for 72 hours. Additionally, a control sample (tap water) was tested. After 3 days, the roots were measured.

2.6 Data analysis

Statistical analysis of all data obtained in the liquid cultures assays was performed through the t-test and the significance levels were set at 95% (p<0.05).

3 RESULTS AND DISCUSSION

3.1 In vitro evaluation of pesticides tolerance and removal by *Pleurotus* fungi

After the incubation period, it was observed mycelial growth of *P. ostreatus* and *P. eryngii* for the treatments of liquid cultures containing bentazon and 2,4-D (T1_{BZ} and T1_{2,4D}). Therefore, the studied fungi may tolerate the pesticides, growing in liquid environments that are similar to the real conditions

of pesticide concentrations.

The biomass dry weight of fungi and the pesticide concentrations after the incubation period for all the studied treatments is presented in Table 1.

Table 1 - Biomass dry weight of fungi and the pesticide concentrations of each treatment after the incubation period.

Treatment	Biomass dry weight (g)	Pesticide concentration (g/L)
CT _{BZ}	0.001 ± 0.000 a	0
T0 _{BZ}	0	0.56 ± 0.03 a
T1 _{BZ}	1.585 ± 0.247 b	0.76 ± 0.01 b
CT _{2,4D}	0.001 ± 0.000 a	0
T0 _{2,4D}	0	0.03 ± 3.72 c
T1 _{2,4D}	0.012 ± 0.002 c	0.03 ± 3.29 c

* vertical lowercase letters indicate the results of comparisons between treatments for each measured variable. Equal letters indicate means that did not statistically differ from each other (t-test, $p \leq 0, 05$).

Treatments T1_{BZ} and T1_{2,4D} presented biomass dry weight statistically different. The fungal biomass of *P. ostreatus* cultivated in a liquid medium containing bentazon was higher than the fungal biomass of *P. eryngii* cultivated in a liquid medium containing 2,4-D, though the cultivation time (7 and 21 days, respectively) as indicated in Table 1. The biomass dry weight of control treatments (CT_{BZ} and CT_{2,4D}) represented just the mycelial disks inoculated in each one due to the absence of an energy source for fungi growth. No biomass dry weight was found for initial treatments (T0_{BZ} and T0_{2,4D}) because of the absence of fungi.

The pesticide concentration of the initial solutions (T0_{BZ} and T0_{2,4D}) was lower than the treatments with fungi growth (T1_{BZ} and T1_{2,4D}). It was observed that bentazon and 2,4-D concentration after the incubation period decreased when compared to the concentration added at the beginning of the experiment (4.5 g L⁻¹ and 5.4 g L⁻¹) independently of the fungi inoculation. This indicates that the herbicide may have been degraded or volatilized during the incubation time.

The results of GC/MS analysis obtained for 2,4-D treatments have no significant statistical difference between them. However, the bentazon concentration was statistically different between the treatments T0_{BZ} and T1_{BZ} as well as about 2,4-D treatments.

In the *in vitro* evaluation of bentazon (4.5 g L⁻¹) and 2,4-D (5.4 g L⁻¹) tolerance by *P. ostreatus* and *P. eryngii*, respectively, the results showed that the fungi can grow up in aqueous media containing these pesticides and, consequently, can tolerate them. The obtained results match those observed in different studies (COELHO-MOREIRA et al., 2018; PEREIRA et al., 2013; FREITAS et al., 2017; GANASH et al., 2016; MARINHO et al., 2017) that reported the fungi tolerance and their potential as agent degradation of pesticides. The biomass dry weight of *P. ostreatus* and *P. eryngii* suggests that the fungi can use bentazon and 2,4-D, respectively, as a source of nutrients. Several reports have shown the ability of *P. ostreatus* to grow in liquid medium with pesticides, such as chlorotoluron, isoproturon, diuron, linuron, propanil, malathion, lindane, and 2,4-dichlorophenol (GANASH et al., 2016; SILVA et al., 2009; RIGAS et al., 2005; KHADRANI et al., 1999; TORRES-DUARTE et al., 2009). Although, some of these experiments highlighted that *P. ostreatus* mycelial growth was inhibited by increasing the concentration of the pesticide (malathion, lindane, and 2,4-dichlorophenol).

Previous studies showed that other basidiomycetes can grow in the presence of bentazon. Coelho-Moreira et al. (2018) studied bentazon removal by *Ganoderma lucidum* in a solid culture medium, using

corn cob as substrate. Despite the authors observed that the herbicide had a negative effect on the mycelial growth of *G. lucidum*, the fungus could grow up in bentazon concentrations below 14.4 g L⁻¹. Castillo et al. (2001) observed the development of *Phanerochaete chrysosporium* in the presence of bentazon (0.22 g L⁻¹) in bioreactors with unsterile wheat straw. In both studies, the fungi could degrade the herbicide.

Silva et al. (2009) observed similar behavior of compound removal by *P. ostreatus* in an experiment where the 2,4-D concentration was 23% lower than the added amount. The degradation of 2,4-D in different concentrations by *T. versicolor* (Hernández Mendieta et al., 2013) and *P. chrysosporium* (DONNELLY et al., 1993) was also reported, though some high concentrations can inhibit their performance (HERNÁNDEZ MENDIETA et al., 2013).

In liquid cultures, the inability of *P. ostreatus* to remove bentazon and *P. eryngii* to remove 2,4-D was observed in this study. Despite the fungal tolerance to pesticides, the cultivation conditions, such as exposure time and pesticide concentration, may not favor the removal of these substances. The same behavior was observed by Donnelly et al. (1993) in an experiment using atrazine as a pollutant, where all fungi grow up in the medium containing the pesticide, but none removed it. In addition, the role of *P. eryngii* and its enzymes have been explored in degrading aromatic hydrocarbon as DDT (PURNOMO; MAULIANAWATI; KAMEI, 2019), fluorene (HADIBARATA; KRISTANTI, 2014), naphthalene (HADIBARATA et al., 2013), and bisphenolic compounds (CHANG; CHANG, 2016).

The degradation of pollutants by fungi is influenced by several factors, including the tested strain, enzyme production, pesticide initial concentration, exposure time, and cultivation conditions (SERBENT et al., 2019). In some organisms, when nutritional sources are present in the medium, the production of enzymes used to metabolize other carbon sources is limited to save energy (MARINHO et al., 2017), which may slow or decrease pesticide removal. Moreover, the mechanisms in the microbial conversion of a pesticide involve secondary metabolism (VALLI; GOLD, 1991), which strongly depends on the cultivation time.

Additionally, the need for supplementation of the culture medium using carbon and nitrogen sources depends on the fungus species. Ganash et al. (2016) observed that the degradation of malathion by *P. ostreatus* was increased when the mineral liquid medium was supplemented with lignin at 0,2%. In that study, the removal of the pesticide reached 76% after 25 days of culture. However, in the presence of *P. ostreatus*, the degradation of the herbicide 2,4-DCP in a liquid medium with wheat straw was better in the absence of glucose, reaching 54% of removal in 96 hours of cultivation (SILVA et al., 2009). According to Donnelly et al. (1993), some fungi needed nitrogen supplementation in the culture media for the complete degradation of 2,4-D. Coelho-Moreira et al. (2018) reported that after 5 days of *G. lucidum* cultivation in a solid and a liquid culture medium with 7.2 g L⁻¹ and 0.6 g L⁻¹ of bentazon respectively, there was no significant removal of the compound. However, after 10 days of treatment, it was detected a removal of 88% of bentazon in the solid medium and 53% in the liquid medium. In experiments in a sucrose liquid medium, Ferreira-Guedes et al. (2012) observed no degradation of 2,4-D by *Penicillium chrysogenum* fungus after the first week of the experiment. However, after 25 days of culture, 18% of this compound was removed. According to these findings, it can be inferred that longer detention time could positively influence the mycoremediation of this herbicide.

Under certain cultivation conditions, basidiomycete fungi can produce nonspecific extracellular enzymes that promote the degradation of pollutant molecules such as laccases, manganese peroxidases, and lignin peroxidases. *P. eryngii* and *P. ostreatus* produce versatile peroxidase that is capable to oxidate aromatic compounds (BANSAL; KANWAR, 2013), such as the ones present in organochloride pesticides. Additionally, Mougín et al. (2000) analyzed the degradation of the herbicide isoxaflutole by the fungi *Phanerochaete chrysosporium* and *Trametes versicolor* in a liquid medium. The authors noted that the metabolization of the herbicide occurred mainly at the end of the growth phase, after 15

days of cultivation of both fungi. This may be related to the production of oxidative enzymes such as laccases. Coelho-Moreira et al. (2018) reported that the herbicides bentazon (20 mM) and diuron (80 μM) stimulated laccase production above 140 and 90 U g^{-1} , respectively, at 10 days of cultivation.

3. 2 Toxicity evaluation of treated water by *Allium* test

The results obtained in the *Allium* test for both bentazon and 2,4-D pesticides with *P. ostreatus* and *P. eryngii*, respectively, are shown in Table 2.

Table 2 - Root growth in each studied treatment in the *Allium* test.

Treatment	Root growth (cm)
CT _{BZ}	4.30 \pm 0.30 a
T0 _{BZ}	0.45 \pm 0.15 b
T1 _{BZ}	0.55 \pm 0.05 b
CT _{2,4D}	3.78 \pm 0.96 a
T0 _{2,4D}	0.33 \pm 0.03 b
T1 _{2,4D}	0.45 \pm 0.13 b

* Vertical lowercase letters indicate the results of comparisons between treatments for each measured variable. Equal letters indicate means that did not statistically differ from each other (t-test, $p \leq 0, 05$).

The samples T0_{BZ}, T1_{BZ}, T0_{2,4D}, and T1_{2,4D} presented low root growth when compared to tap water (CT_{BZ} and CT_{2,4D}) (Table 2). The tap water samples presented a significant difference compared to the samples in that the roots were exposed to treatments containing bentazon and 2,4-D pesticides. All the treatments containing pesticides present no significant statistical difference in root growth.

Many tests have been developed to identify the harmful effects of chemicals on animals and plants, with emphasis on plant-based assays for being short-term and low-cost alternatives (YILMAZ et al., 2018). To verify the toxicity of samples, some assays can be performed, such as the *Allium* test. This is an interesting tool for conducting toxicity and mutagenicity research using onions (FREITAS et al., 2017; MARINHO et al., 2017) once they are cultivated during all seasons and have a low economic cost. Its roots are activated when the bulb contacts water, however, the presence of toxic compounds in the water reduces the root growth due to the delay in division cellular (MARINHO et al., 2017).

A. cepa test is a useful bioindicator of cytotoxicity and genotoxicity and serves as an alert for the population that uses pesticides indiscriminately (DATTA et al., 2018). The results obtained in the *Allium cepa* test highlighted the toxicity of the liquid culture with bentazon and 2,4-D due to the low root growth of onions observed in the samples containing the pesticides when compared to tap water. The toxicity of the samples remains after the treatments with *P. ostreatus* and *P. eryngii* because of the non-removal of pesticide compounds in the studied cultivation conditions, as demonstrated in GC/MS analysis.

No data was found on toxicity tests with bentazon in the literature. However, other studies demonstrate the detoxification of pesticides by basidiomycete fungi. In *Lactuca sativa* bioassays, Coelho-Moreira et al. (2018) observed a reduction in toxicity after fungal treatment compared to untreated samples. In another study, the *Allium cepa* test showed an inhibition decrease from 100% to 54% after atrazine treatment with *Aspergillus niger* in liquid culture after the incubation period of 7 days (MARINHO et al., 2017). Additionally, these authors obtained favorable results in the removal of pesticides methyl parathion and atrazine. After seven days of treatment with the fungus *Aspergillus niger*, the *Allium cepa* test demonstrated that the toxicity of the treated sample had decreased, as there was a growth of the

roots of the onions.

Fatima and Ahmad (2006) analyzed the EROD (o ethoxy resorufin O-demethylase) activity in *Allium cepa* exposed to 2,4-D, as a response of the organism to the presence of pollutants in the aquatic environment. The 2,4-D caused about a 40-fold rise in EROD at a dose of 1.2 ppb. The authors concluded that the strategy allows the detection of this pesticide in water monitoring studies before using analytical techniques like HPLC.

Özkul et al. (2016) investigated the cytogenetic effects of different concentrations of 2,4-D (0.67, 1.34, 2.01, 2.68, 3.35, and 4.02 mg/L) on *Allium cepa* bulblets' root tips treated for 24 and 48 h. The results showed that the herbicide affected mitotic index (MI), whose value increased significantly at three lower concentrations (0.67, 1.34, 2.01 mg/L) after treatment with 2,4-D for 24 h and decreased significantly at higher concentrations (2.68, 3.35, and 4.02 mg/L). Also, the authors showed that roots were very sensitive to the herbicide if they had been in touch with 2,4-D (4.02 mg/L) for a longer time (48 h).

The 2,4-D incidence and the cytotoxic and genotoxic potential of sediment elutriate of rivers from southern Brazil, by *Allium cepa* test, as evidenced by Rambo et al. (2017) while Martins and Campos Pereira (2018) reported that the herbicide Tordon®, which has in its formulation the compound 4-amino-3,5,6-trichloro picolinic acid (Picloram - 103,6 g/L) and 2,4-D (402 g/L) inhibited (concentrations of 0.1%, 1.0%, 10%, and 100% of the herbicide) or significantly decreased the length of the onion roots at concentrations of 0.01%, 0.001%, and 0.0001%.

Besides the non-removal of bentazon and 2,4-D, the toxicity results obtained in our study may be explained by the possible occurrence of pesticide metabolites that are not metabolized by the fungi and inhibit the onions-root growth.

There are different mechanisms involved in pesticide conversion by microorganisms. Depending on the environmental conditions, each mechanism can produce metabolites that are different from the initial compound (SERBENT et al., 2019). Consequently, even the pesticide is susceptible to degradation by fungi, in some cases, its degradation process can produce compounds that are more toxic than the pesticide itself (LURQUIN, 2016). On the other hand, some metabolites may be more accessible to fungal degradation compared to the initial compound, as proved by Vroumsia et al. (2005). The authors showed that different fungal strains more efficiently degraded the 2,4-DCP than the 2,4-D, suggesting that this behavior can occur because of the compounds' bindings and free phenolic radicals. Besides, other fungal strains better-degraded 2,4-D than 2,4-DCP, leading to hypothesizing that the conversion of 2,4-D and 2,4-DCP are catalyzed by different enzymes (VROUMSIA et al. 2005). Similarly, the metabolites of bentazon on surface water were identified as more toxic than the initial substance for fish and invertebrates while the opposite occurs for other aquatic organisms (EFSA, 2015).

Considering the obtained data on this study it may be hypothesized that the studied fungi are capable to tolerate 2,4-D and bentazon, but the cultivation conditions were not favorable to the degradation of the compounds. This can be explained by the fact that the degradation ability depends on the fungi species, the contaminant and its concentration, and the nitrogen concentration in the culture media (DONNELLY et al., 1993). In addition, these authors also concluded that it may have an increase in degradation rate with the biomass increase and the enzymatic activity when there is a high amount of carbohydrate in the culture media. Finally, the basidiomycete fungi are considered potential accumulators (BOSCO; MOLLEA, 2019). The aromatic herbicides' degradation appears to be through the compounds' incorporation into the fungus tissue and not through the complete degradation of the contaminant (mineralization) (DONNELLY et al., 1993).

Thus, the use of *P. ostreatus* and *P. eryngii* for mycoremediation of bentazon and 2,4-D in contaminated effluents needs to be further studied. To investigate the role of this organism in the treatment of bentazon and 2,4-D contaminated effluents, new experimental conditions must be tested. In addition, the tests

should include the monitoring and quantification of enzymatic activity as well as the pesticide degradation metabolites. As a result, it could be possible to know the moment that *P. ostreatus* and *P. eryngii* start to release oxidative enzymes and if it contributes to bentazon and 2,4-D metabolization and its metabolites.

Also, the fungi tolerance to herbicides should investigate the mycelium characteristics after herbicide exposition regarding reproductive structures, hyphae feature, and exposition time–structure-property relationships, which are relevant to a variety of applications of mycelium as is the case of a bioreactor for the treatment of contaminated effluents (SERBENT et al., 2020).

4 CONCLUSION

Due to the characteristics of WRF, the purpose of the present study was to evaluate the potential use of Basidiomycota fungi as degradation agents of bentazon and 2,4-D by assessing their tolerance, growth, and removal capacity of pesticides in liquid cultures. In summary, the fungus *P. ostreatus* and *P. eryngii*, grew up in liquid culture mediums with 4.5 g L⁻¹ of bentazon and 5.4 g L⁻¹ of 2,4-D, respectively, and tolerate this environment. Despite that, the fungi were not efficient in removing the pesticides after seven and twenty-one days of cultivation. The *Allium cepa* test evidenced that samples containing pesticides were still toxic after the treatments with fungi, in the studied conditions. The results represent the starting point of mycoremediation processes for future applications on a pilot and full scale.

Disclosure statement

There are no relevant financial or non-financial competing interests to report. No potential competing interest was reported by the authors.

Acknowledgments

The authors thank the financial support provided by Santa Catarina State University. Also, the authors are thankful to the Federal University of Santa Catarina, especially to the laboratory of microorganisms and biotechnological processes, for providing the strain of *Pleurotus ostreatus* and *Pleurotus eryngii*; and to the Coordination for the Improvement of Higher Education Personnel – Brazil (CAPES).

5 REFERENCES

- AKHTAR, N.; MANNAN, M. A. Mycoremediation: Expunging environmental pollutants. **Biotechnology Reports**, v. 26, p. e00452, 2020. DOI: <https://doi.org/10.1016/j.btre.2020.e00452>
- BANSAL, N.; KANWAR, S. S. Peroxidase(s) in environment protection. **The Scientific World Journal**, v. 2013, p. 1-9, 2013. DOI: <https://doi.org/10.1155/2013/714639>
- BOSCO, F.; MOLLEA, C. Mycoremediation in Soil. In: SALDARRIAGA-NOREÑA, H.; MURILLO-TOVAR, M. A.; FAROOQ, R.; DONGRE, R.; RIAZ, S. (Eds.). **Environmental Chemistry and Recent Pollution Control Approaches**. London: IntechOpen, 2019, p. 1-17.
- CALDAS, S. S.; BOLZAN, C. M.; GUILHERME, J. R.; SILVEIRA, M. A. K.; ESCARRONE, A. L. V.; PRIMEL, E. G. Determination of pharmaceuticals, personal care products, and pesticides in surface and

treated waters: Method development and survey. **Environmental Science and Pollution Research**, v. 20, n. 8, p. 5855-5863, 2013. DOI: <https://doi.org/10.1007/s11356-013-1650-9>

CASTILLO, M. D. P.; ANDERSSON, A.; ANDER, P.; STENSTRÖM, J.; TORSTENSSON, L. Establishment of the white rot fungus *Phanerochaete chrysosporium* on unsterile straw in solid substrate fermentation systems intended for degradation of pesticides. **World Journal of Microbiology and Biotechnology**, v. 17, p. 627-633, 2001. DOI: <https://doi.org/10.1023/A:1012420422111>

CHANG, B. V.; CHANG, Y. M. Biodegradation of toxic chemicals by *Pleurotus eryngii* in submerged fermentation and solid-state fermentation. **Journal of Microbiology, Immunology and Infection**, v. 49, n. 2, p. 175-181, 2016. DOI: <https://doi.org/10.1016/j.jmii.2014.04.012>

CHEN, Y.; YU, K.; HASSAN, M.; XU, C.; ZHANG, B.; GIN, K. Y-H.; HE, Y. Occurrence, distribution and risk assessment of pesticides in a river-reservoir system. **Ecotoxicology and Environmental Safety**, v. 166, p. 320-327, 2018. DOI: [10.1016/j.ecoenv.2018.09.107](https://doi.org/10.1016/j.ecoenv.2018.09.107)

COELHO-MOREIRA, J. da S.; BRUGNARI, T.; SÁ-NAKANISHI, A. B.; CASTOLDI, R.; SOUZA, C. G. M. de; BRACHT, A.; PERALTA, R. M. Evaluation of diuron tolerance and biotransformation by the white-rot fungus *Ganoderma lucidum*. **Fungal Biology**, v. 122, n. 6, p. 471-478, 2018. DOI: <https://doi.org/10.1016/j.funbio.2017.10.008>

CUPUL, W. C.; ABARCA, G. H.; VÁZQUEZ, R. R.; SALMONES, D.; HERNÁNDEZ, R. G.; GUTIÉRREZ, E. A. Response of ligninolytic macrofungi to the herbicide atrazine: dose-response bioassays. **Revista Argentina de Microbiología**, v. 46, n. 4, p. 348-357, 2014. DOI: [https://doi.org/10.1016/S0325-7541\(14\)70094-X](https://doi.org/10.1016/S0325-7541(14)70094-X)

DATTA, S.; SINGH, J.; SINGH, J.; SINGH, S.; SINGH, S. Assessment of genotoxic effects of pesticide and vermicompost treated soil with *Allium cepa* test. **Sustainable Environment Research**, v. 28, p. 171-178, 2018. DOI: <https://doi.org/10.1016/j.serj.2018.01.005>

CASTRO LIMA, J. A. M. de; LIMA, J. A. M. de C.; LABANOWSKI, J.; BASTOS, M. C.; ZANELLA, R.; PRESTES, O. D.; VARGAS, J. P. R. de; MONDAMERT, L.; GRANADO, E.; TIECHER, T.; ZAFAR, M.; TROIAN, A.; LE GUET, T.; SANTOS, D. R. dos. "Modern agriculture" transfers many pesticides to watercourses: a case study of a representative rural catchment of southern Brazil. **Environmental Science and Pollution Research**, v. 27, p. 10581-10598, 2020. DOI: <https://doi.org/10.1007/s11356-019-06550-8>

SOUZA, D. F. de; TYCHANOWICZ, G. K.; SOUZA, C. G. M. de; PERALTA, R. M. Co-production of ligninolytic enzymes by *Pleurotus pulmonarius* on wheat bran solid state cultures. **Journal of Basic Microbiology**, v. 46, n. 2, p. 126-134, 2006. DOI: <https://doi.org/10.1002/jobm.200510014>

DONNELLY, P. K.; ENTRY, J. A.; CRAWFORD, D. L. Degradation of atrazine and 2,4-dichlorophenoxyacetic acid by mycorrhizal fungi at three nitrogen concentrations in vitro. **Applied and Environmental Microbiology**, v. 59, n. 8, p. 2642-2647, 1993. DOI: <https://doi.org/10.1128/aem.59.8.2642-2647.1993>

ECHEVERRI GONZÁLEZ, A. F.; ZÚÑIGA-BENÍTEZ, H.; PEÑUELA, G. A. Removal of herbicide 2,4-D using constructed wetlands at pilot scale. **Emerging Contaminants**, v. 5, p. 303-307, 2019. DOI: <https://doi.org/10.1016/j.emcon.2019.04.001>

doi.org/10.1016/j.emcon.2019.09.001

EUROPEAN FOOD SAFETY AUTHORITY (EFSA). Conclusion on the peer review of the pesticide risk assessment of the active substance bentazone. **European Food Safety Authority Journal**, v. 13, n. 4, p. 4077, 2015. DOI: <https://doi.org/10.2903/j.efsa.2015.4077>

FATIMA, R. A.; AHMAD, M. *Allium cepa* derived EROD as a potential biomarker for the presence of certain pesticides in water. **Chemosphere**, v. 62, n. 4, p. 527-537, 2006. DOI: <https://doi.org/10.1016/j.chemosphere.2005.06.032>

FERREIRA-GUEDES, S.; MENDES, B.; LEITÃO, A. L. Degradation of 2,4-dichlorophenoxyacetic acid by a halotolerant strain of *Penicillium chrysogenum*: Antibiotic production. **Environmental Technology**, v. 33, p. 677-686, 2012. DOI: <https://doi.org/10.1080/09593330.2011.588251>

FREITAS, L. M. C.; BARBOSA, B. C. A.; RODRIGUES, K.; MARINHO, G. Employment of *Aspergillus Niger* AN 400 in batch reactors to remove pesticide aqueous matrix. **Engenharia Sanitaria e Ambiental**, v. 22, n. 6, p. 1175-1185, 2017. DOI: <https://doi.org/10.1590/S1413-41522017138266>

GANASH, M. A.; ABDEL GHANY, T. M.; REYAD, A. M. *Pleurotus ostreatus* as a biodegradator for organophosphorus insecticide malathion. **Journal of Environmental & Analytical Toxicology**, v. 6, n. 3, p. 369, 2016. DOI: <https://doi.org/10.4172/2161-0525.1000369>

HADIBARATA, T.; TEH, Z. C.; RUBIYATNO; ZUBIR, M. M. F. A.; KHUDHAIR, A. B.; YUSOFF, A. R. M.; SALIM, M. R.; HIDAYAT, T. Identification of naphthalene metabolism by white rot fungus *Pleurotus eryngii*. **Bioprocess and Biosystems Engineering**, v. 36, n. 10, p. 1455-1461, 2013. DOI: <https://doi.org/10.1007/s00449-013-0884-8>

HADIBARATA, T.; KRISTANTI, R. A. Potential of a white-rot fungus *Pleurotus eryngii* F032 for degradation and transformation of fluorene. **Fungal Biology**, v. 118, n. 2, p. 222-227, 2014. DOI: <https://doi.org/10.1016/j.funbio.2013.11.013>

HARMS, H.; SCHLOSSER, D.; WICK, L. Y. Untapped potential: Exploiting fungi in bioremediation of hazardous chemicals. **Nature Reviews Microbiology**, v. 9, p. 177-192, 2011. DOI: <https://doi.org/10.1038/nrmicro2519>

HERNÁNDEZ MENDIETA, E.; SÁNCHEZ, D. G.; MARTÍNEZ, V.L.; TEJACAL, I. A.; RODRÍGUEZ, M. A.; TORRES O. G. V.; FERNÁNDEZ, E. M.; LARA, M. H.; MIRANDA, A. S. Identificación del agente causal de la pudrición blanca en Morelos, México. **Revista Colombiana de Biotecnología**, v. 15, n. 2, p. 178-185, 2013. DOI: <https://doi.org/10.15446/rev.colomb.biote.v15n2.41744>

KAPAH, M.; SACHDEVA, S. Mycoremediation potential of *Pleurotus* species for heavy metals: a review. **Bioresources and Bioprocessing**, v. 4, p. 32, 2017. DOI: <https://doi.org/10.1186/s40643-017-0162-8>

KHADRANI, A.; SEIGLE-MURANDI, F.; STEIMAN, R.; VROUMSIA, T. Degradation of three phenylurea herbicides (chlortoluron, isoproturon and diuron) by micromycetes isolated from soil. **Chemosphere**, v. 38, n. 13, p. 3041-3050, 1999. DOI: [https://doi.org/10.1016/S0045-6535\(98\)00510-4](https://doi.org/10.1016/S0045-6535(98)00510-4)

KIST, C. P.; SCHERER, C. E.; SOARES, M.; RODRIGUES, M.B. Biodegradation of nitroaromatic compounds in red water by white rot fungi *Pleurotus ostreatus* and *floridae*. **Ambiente & Água**, v. 15, n. 6, p. e2594, 2020. <https://doi.org/10.4136/ambi-agua.2594>

KÖCK-SCHULMEYER, M.; GINEBREDÀ, A.; POSTIGO, C.; GARRIDO, T.; FRAILE, J.; ALDA, M. L. de; BARCELÓ, D. Four-year advanced monitoring program of polar pesticides in groundwater of Catalonia (NE-Spain). **Science of the Total Environment**, v. 470-471, p. 1087-1098, 2014. DOI: <https://doi.org/10.1016/j.scitotenv.2013.10.079>

LEONHARDT, S.; HOPPE, B.; STENGEL, E.; NOLL, L.; MOLL, J.; BÄSSLER, C.; DAHL, A.; BUSCOT, F.; HOFRICHTER, M.; KELLNER, H. Molecular fungal community and its decomposition activity in sapwood and heartwood of 13 temperate European tree species. **PLoS One**, v. 14, n. 2, p. e0212120, 2019. DOI: <https://doi.org/10.1371/journal.pone.0212120>

LURQUIN, P. F. Production of a toxic metabolite in 2,4-D-resistant GM crop plants. **3 Biotech**, v. 6, n. 1, p. 82, 2016. DOI: <https://doi.org/10.1007/s13205-016-0387-9>

MARINHO, G.; BARBOSA, B. C. A.; RODRIGUES, K.; AQUINO, M.; PEREIRA, L. Potential of the filamentous fungus *Aspergillus niger* AN 400 to degrade Atrazine in wastewaters. **Biocatalysis and Agricultural Biotechnology**, v. 9, p. 162-167, 2017. DOI: <https://doi.org/10.1016/j.bcab.2016.12.013>

MARTINS, H.; CAMPOS PEREIRA, F. D. Avaliação dos efeitos tóxicos do agroquímico Tordon® sobre os organismos teste *Lactuca sativa* e *Allium cepa*. **Visão Acadêmica**, v. 19, n. 2, p. 55-69, 2018. DOI: <http://dx.doi.org/10.5380/acd.v19i2.59763>

MOUGIN, C.; BOYER, F. D.; CAMINADE, E.; RAMA, R. Cleavage of the diketone nitrile derivative of the herbicide isoxaflutole by extracellular fungal oxidases. **Journal of Agricultural and Food Chemistry**, v. 48, n. 10, p. 4529-4534, 2000. DOI: <http://dx.doi.org/10.1021/jf000397q>

ÖZKUL, M.; ÖZEL, Ç. A.; YÜZBAŞIOĞLU, D.; ÜNAL, F. Does 2,4-dichlorophenoxyacetic acid (2,4-D) induce genotoxic effects in tissue cultured *Allium* roots? **Cytotechnology**, v. 68, p. 2395-2405, 2016. DOI: <http://dx.doi.org/10.1007/s10616-016-9956-3>

PEREIRA, P. M.; TEIXEIRA, R. S. S.; OLIVEIRA, M. A. L. de; SILVA, M. da; LEITÃO, V. S. F. Optimized atrazine degradation by *Pleurotus ostreatus* INCQS 40310: an alternative for impact reduction of herbicides used in sugarcane crops. **Journal of Microbial and Biochemical Technology**, v. S12, p. 006, 2013. DOI: <http://dx.doi.org/10.4172/1948-5948.S12-006>

POSTIGO, C.; ALDA, M. L. S. de; BARCELÓ, D.; GINEBREDÀ, A.; GARRIDO, T.; FRAILE, J. Analysis and occurrence of selected medium to highly polar pesticides in groundwater of Catalonia (NE Spain): An approach based on on-line solid phase extraction-liquid chromatography-electrospray-tandem mass spectrometry detection. **Journal of Hydrology**, v. 383, n. 1-2, p. 83-92, 2010. DOI: <https://doi.org/10.1016/j.jhydrol.2009.07.036>

PURNOMO, A. S.; MORI, T.; KAMEI, I.; NISHII, T.; KONDO, R. Application of mushroom waste medium from *Pleurotus ostreatus* for bioremediation of DDT-contaminated soil. **International Biodeterioration and Biodegradation**, v. 64, n. 5, p. 397-402, 2010. DOI: <https://doi.org/10.1016/j.ibiod.2010.04.007>

PURNOMO, A. S.; MAULIANAWATI, D.; KAMEI, I. 2019. *Ralstonia pickettii* enhance the DDT biodegradation by *Pleurotus eryngii*. **Journal of Microbiology and Biotechnology**, v. 29, n. 9, p. 1424-1433. DOI: <https://doi.org/10.4014/jmb.1906.06030>

QUINTANA, J.; DE LA CAL, A.; BOLEDA, M. R. Monitoring the complex occurrence of pesticides in the Llobregat basin, natural and drinking waters in Barcelona metropolitan area (Catalonia, NE Spain) by a validated multi-residue online analytical method. **Science of the Total Environment**, v. 692, p. 952-965, 2019. DOI: <https://doi.org/10.1016/j.scitotenv.2019.07.317>

RAMBO, C. L. ZANOTELLI, P.; DALEGRAVE, D.; NEZ D. de; SZCZEPANIK, J.; CARAZEK, F.; FRANSCESCON, F.; ROSEMBERG, D. B.; SIEBEL, A. M.; DAL MAGRO, J. Hydropower reservoirs: cytotoxic and genotoxic assessment using the *Allium cepa* root model. **Environmental Science and Pollution Research**, v. 24, p. 8759-8768, 2017. DOI: <https://doi.org/10.1007/s11356-017-8509-4>

RIGAS, F.; DRITSA, V.; MARCHANT, R.; PAPADOPOULOU, K.; AVRAMIDES, E.J.; HATZIANESTIS, I. Biodegradation of lindane by *Pleurotus ostreatus* via central composite design. **Environment International**, v. 31, n. 2, p. 191-196, 2005. DOI: <https://doi.org/10.1016/j.envint.2004.09.024>

RILEY, R.; SALAMOV, A. A.; BROWN, D. W.; GRIGORIEV, I. V. Extensive sampling of basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood decay fungi. **Proceedings of the National Academy of Sciences of the United States of America**, v. 111, n. 27, p. 9923-9928, 2014. DOI: <https://doi.org/10.1073/pnas.1400592111>

SALMAN, J. M.; HAMEED, B. H. Effect of preparation conditions of oil palm fronds activated carbon on adsorption of bentazon from aqueous solutions. **Journal of Hazardous Material**, v. 175, n. 1-3, p. 133-137, 2010. DOI: <https://doi.org/10.1016/j.jhazmat.2009.09.139>

SANTOS, E. C. et al. 2019. Degradação do repelente DEET pelas lacases do *Pleurotus ostreatus*. **Revista de Divulgação Científica Sena Aires**, 8(2):160-169.

SERBENT, M. P.; REBELO, A. M.; PINHEIRO, A.; GIONGO, A.; TAVARES, L. B. B. Biological agents for 2,4-dichlorophenoxyacetic acid herbicide degradation. **Applied Microbiology and Biotechnology**, v. 103, p. 5065-5078, 2019. DOI: <https://doi.org/10.1007/s00253-019-09838-4>

SERBENT, M. P.; GUIMARÃES, D. K. S.; DRECHSLER-SANTOS, E. R.; HELM, C. V.; GIONGO, A.; TAVARES, L. B. B. Growth, enzymatic production and morphology of the white-rot fungi *Lentinus crinitus* (L.) Fr. upon 2,4-D herbicide exposition. **International Journal of Environmental Science and Technology**, v. 17, p. 2995-3012, 2020. DOI: <https://doi.org/10.1007/s13762-020-02693-1>

SJERPS, R. M. A.; KOOIJ, P. J. F.; VAN LOON, A.; VAN WEZEL, A.P. Occurrence of pesticides in Dutch drinking water sources. **Chemosphere**, v. 235, p. 510-518, 2019. DOI: <https://doi.org/10.1016/j.chemosphere.2019.06.207>

SILVA, H. H. B. da; SCHNEIDER, A. L. S.; WISBECK, E.; FURLAN, S. A. Biodegradation of 2,4 dichlorophenol by *Pleurotus ostreatus* DSM 1833. **Brazilian Archives of Biology and Technology**, v. 52, n. 6, p. 1563-1570, 2009. DOI: <https://doi.org/10.1590/S1516-89132009000600028>

SINGH, H. **Mycoremediation**: Fungal Bioremediation. 1. ed. Hoboken: John Wiley & Sons, 2006, 616 p.

TORRES-DUARTE, C.; ROMAN, R.; TINOCO, R.; VAZQUEZ-DUHALT, R. Halogenated pesticide transformation by a laccase-mediator system. **Chemosphere**, v. 77, n. 5, p. 687-692, 2009. DOI: <https://doi.org/10.1016/j.chemosphere.2009.07.039>

US ENVIRONMENTAL PROTECTION AGENCY (EPA). **Method 8270D (SW846)**: Semivolatile Organic Compounds by Gas Chromatography/Mass Spectrometry. Washington, DC: EPA, 2014.

VALDEZ-VAZQUEZ, I.; ROBLEDO-RIZO, J. G.; MUÑOZ-PÁEZ, K. M.; PÉREZ-RANGEL, M.; RUIZ-AGUILAR, G. M. L. Simultaneous hydrogen production and decolorization of denim textile wastewater: kinetics of decolorizing of indigo dye by bacterial and fungal strains. **Brazilian Journal of Microbiology**, v. 51, p. 701-709, 2020. DOI: <https://doi.org/10.1007/s42770-019-00157-4>

VALLI, K.; GOLD, M. H. Degradation of 2,4-dichlorophenol by the lignin-degrading fungus *Phanerochaete chrysosporium*. **Journal of Bacteriology**, v. 173, n. 1, p. 345-352, 1991. DOI: <https://doi.org/10.1128/jb.173.1.345-352.1991>

VIEIRA, D. C.; NOLDIN, J. A.; DESCHAMPS, F. C.; RESGALLA JR, C. Ecological risk analysis of pesticides used on irrigated rice crops in southern Brazil. **Chemosphere**, v. 162, p. 48-54, 2016. DOI: <https://doi.org/10.1016/j.chemosphere.2016.07.046>

VROUMSIA, T.; STEIMAN, R.; SEIGLE-MURANDI, F.; BENOIT-GUYOD, J-L. Fungal bioconversion of 2,4-dichlorophenoxyacetic acid (2,4-D) and 2,4-dichlorophenol (2,4-DCP). **Chemosphere**, v. 60, n. 10, p. 1471-1480, 2005. <https://doi.org/10.1016/j.chemosphere.2004.11.102>

XIE, H.; WANG, X.; CHEN, J.; LI, X.; JIA, G.; ZOU, Y.; ZHANG, Y.; CUI, Y. Occurrence, distribution and ecological risks of antibiotics and pesticides in coastal waters around Liaodong Peninsula, China. **Science of the Total Environment**, v. 656, p. 946-951, 2019. DOI: <https://doi.org/10.1016/j.scitotenv.2018.11.449>

XU, M.; HUANG, H.; LI, N.; LI, F.; WANG, D.; LUO, Q. Occurrence and ecological risk of pharmaceuticals and personal care products (PPCPs) and pesticides in typical surface watersheds, China. **Ecotoxicology and Environmental Safety**, v. 175, p. 289-298, 2019. DOI: <https://doi.org/10.1016/j.ecoenv.2019.01.131>

YILMAZ, S.; MARAKLI, S.; YUZBASIOGLU, G.; GOZUKIRMIZI, N. Short-term mutagenicity test by using IRAP molecular marker in rice grown under herbicide treatment. **Biotechnology & Biotechnological Equipment**, v. 32, n. 4, p. 923-928, 2018. DOI: <https://doi.org/10.1080/13102818.2018.1474137>

YADAV, I. C.; DEVI, N. L.; SYED, J. H.; CHENG, Z.; LI, J.; ZHANG, G.; JONES, K. C. Current status of persistent organic pesticides residues in air, water, and soil, and their possible effect on neighboring countries: A comprehensive review of India. **Science of the Total Environment**, v. 511, p. 123-137, 2015. DOI: <https://doi.org/10.1016/j.scitotenv.2014.12.041>

ZAPANA-HUARACHE, S. V.; ROMERO-SÁNCHEZ, C. K.; DUEÑAS GONZA, A. P.; TORRES-HUACO, F. D.; LAZARTE RIVERA, A. M. Chromium (VI) bioremediation potential of filamentous fungi isolated from Peruvian tannery industry effluents. **Brazilian Journal of Microbiology**, v. 51, p. 271-278, 2020. DOI: <https://doi.org/10.1007/s42770-019-00209-9>

ZUANAZZI, N. R.; GHISI, N. C.; OLIVEIRA, E. C. Analysis of global trends and gaps for studies about 2,4-D herbicide toxicity: A scientometric review. **Chemosphere**, v. 241, p. 125016, 2020. DOI: <https://doi.org/10.1016/j.chemosphere.2019.125016>